

AD-A186 425

ON HYPERCONTRACTIVITY OF ALPHA-STABLE RANDOM VARIABLES 1/1

0 ALPHA 2(U) NORTH CAROLINA UNIV AT CHAPEL HILL DEPT OF
STATISTICS J SZULGA JUL 87 TR-196 AFOSR-TR-87-1121


UNCLASSIFIED

F49620-85-C-0144

F/G 12/3

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A186 425

DTIC FILE COPY ①

REPORT DOCUMENTATION PAGE


1a. REPORT SECURITY CLASSIFICATION			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			Approved for public release; distribution unlimited.		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 196			5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR. 87-1121		
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina		6b. OFFICE SYMBOL (if applicable)		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6c. ADDRESS (City, State, and ZIP Code) 321-A Phillips Hall 039-A Statistics Dept. Chapel Hill, NC 27514		7b. ADDRESS (City, State, and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION AFOSR		8b. OFFICE SYMBOL (if applicable) NM		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85C 0144.	
8c. ADDRESS (City, State, and ZIP Code) AFOSR/NM Bldg 410 Bolling AFB DC 20332-6448		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO. 61102F		PROJECT NO. 2304	
				TASK NO. A-5	
				WORK UNIT ACCESSION NO.	
11. TITLE (Include Security Classification) On Hypercontractivity of α -stable Random Variables, $0 < \alpha < 2$					
12. PERSONAL AUTHOR(S) Dr. Jerzy Szulga					
13a. TYPE OF REPORT preprint		13b. TIME COVERED FROM 9/86 TO 8/87		14. DATE OF REPORT (Year, Month, Day) July 1987	
15. PAGE COUNT 13					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	Key Words and Phrases: hypercontractivity, stable random variables, domain of normal attraction.		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>Let θ be a symmetric α-stable random variable, $0 < \alpha < 2$. We show that for any p and q, $0 < q < p < \alpha$, there is a constant $s = s(p, q, \alpha)$ such that</p> $E x + s\theta y ^p)^{1/p} \leq (E x + \theta y ^q)^{1/q}$ <p>for all x and y from a normed space.</p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Maj. Brian Woodruff			22b. TELEPHONE (Include Area Code) (202) 767-5026		22c. OFFICE SYMBOL NM

DTIC
ELECTE
S OCT 06 1987 D
E

AFOSR-TR- 87-1121

CENTER FOR STOCHASTIC PROCESSES

Department of Statistics
University of North Carolina
Chapel Hill, North Carolina



On Hypercontractivity of α -stable Random Variables, $0 < \alpha < 2$

Jerzy Szulga

Technical Report No. 196

July 1987

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

ON HYPERCONTRACTIVITY OF α -STABLE RANDOM VARIABLES, $0 < \alpha < 2$.*)

by Jerzy Szulga^{^)}

Center for Stochastic Processes
University of North Carolina
Chapel Hill, NC 27514

Let θ be a symmetric α -stable random variable, $0 < \alpha < 2$. We show that for any p and q , $0 < q < p < \alpha$, there is a constant $s = s(p, q, \alpha)$ such that

$$E||x + s\theta y||^p)^{1/p} \leq (E||x + \theta y||^q)^{1/q}$$

for all x and y from a normed space.

Key words and phrases: hypercontractivity, stable random variables, domain of normal attraction

AMS 1980 Subject classification: Primary: 60B11, 42C15;

Secondary: 60C42, 46E30, 42C05.

*) A part of the research was conducted during the author's stay at Case Western Reserve University, Cleveland, Ohio. Research partially sponsored by Air Force Office of Scientific Research Grant No. F49620 85C 0144.

^{^)} On leave from Wroclaw University, Poland

1. Introduction

The classical Nelson inequality [9]

$$(1.1) \quad ((|x+s_{p,q}y|^p + |x-s_{p,q}y|^p)/2)^{1/p} \leq ((|x+y|^p + |x-y|^p)/2)^{1/p},$$

where $1 < q \leq p < \infty$, $x, y \in \mathbb{R}$, and $s_{p,q} = ((q-1)/(p-1))^{1/2}$, and its extensions play a fundamental role in the theory of hypercontractive operators [1,4,12], convolution inequalities [11], logarithmic Sobolev inequalities [4], stochastic Ising models [5] and related subjects in harmonic analysis, statistical mechanics and quantum physics, to name just a few areas of great importance nowadays.

Some more sophisticated counterparts of Nelson's inequality were recently applied in the intensively developing theory of multiple stochastic integrals, random multilinear forms and stochastic chaoses, topics originated with Wiener [13] in the late 30's. Using some properties of hypercontractive operators, Borell [2] showed that all p^{th} norms, $p \geq 2$, of Hilbert space valued polynomial chaoses in independent random variables are comparable, generalizing a *fortiori* classical results of Marcinkiewicz, Paley and Zygmund [8,10].

Furthermore, this concept became a basic part of a construction of a stable multiple stochastic integral due to Krakowiak and Szulga [7]. A notion of hypercontractive random variables was introduced for this purpose, having also an intrinsic interest.

Following the key feature of the Nelson inequality, a random variable θ is called hypercontractive with indices $p, q > 0$ in a normed space X if there exists a constant s such that

$$(1.2) \quad (E\{|x + s\theta y|^p\})^{1/p} \leq (E\{|x + \theta y|^q\})^{1/q}$$

for all $x, y \in X$; $\theta \in \text{HC}(p, q, X; s)$ in short.

For example, Borell's extension of (1.1) ([2]) says that for a Bernoulli random variable σ taking the value $+1$ or -1 with probability $\frac{1}{2}$, $\sigma \in \text{HC}(p, q, X; s_{p,q})$ in any normed space X , where $1 < q < p < \infty$.

By a central limit theorem argument, a symmetric Gaussian random variable $\tau \in \text{HC}(p, q, X; s_{p,q})$ with the same parameters p, q and $s_{p,q}$ (cf. also Nelson [9]).

The number $s_{p,q}$ is the best possible constant for τ and σ . Moreover, q must be greater than 1. This is obvious in case of a Bernoulli random variable because, for $t > 0$,

$$E|1 + t\sigma|^q \begin{cases} \leq 1 & \text{if } q \leq 1, \\ > 1 & \text{if } q \geq 1. \end{cases}$$

In case of a Gaussian random variable τ we have, for $0 < q \leq 1 < p$ and $|t| < 1$,

$$E|1 + t\tau|^q \leq 1 + q(q-1)/2 E\tau^2 I\{|\tau| \leq 1\} + \text{const } e^{-1/t},$$

$$E|1 + t\tau|^p \geq 1 + p(p-1)/2 E\tau^2 I\{|\tau| \leq 1\} t^2.$$

The latter inequalities are due to S. Kwapien.

An α -stable symmetric random variable is hypercontractive in an arbitrary normed space if $1 < q < p < \alpha \leq 2$ (Krakowiak and Szulga [6]). Although this is an attribute of a more general class of probability distributions (*ibidem*) integrability and limitation to parameters $q > 1$ seemed like the prerequisite conditions.

We show in this paper that α -stable symmetric random variables are hypercontractive for all α, p, q , with $0 < q < p < \alpha < 2$, in any normed space.

2. Properties of hypercontractive random variables

For a $p > 0$ and a Banach space X we denote by $L_p(X)$ the Banach (Fréchet, if $p < 1$) space of X -valued random variables θ such that $\|\theta\|_p = (E\|\theta\|^p)^{1/p} < \infty$. A function $F = F_k: N^k \rightarrow X$, $k \geq 1$, is called *tetrahedral* if F vanishes outside the "tetrahedron" $D_k = \{\underline{i}_k = (i_1, \dots, i_k): 1 \leq i_1 < \dots < i_k\}$. We define a homogeneous polynomial of degree k on R^N by

$$\langle F_k; (\underline{t})_k \rangle = \sum_{\underline{i}_k \in D_k} F(\underline{i}_k) t_{i_1} \dots t_{i_k}, \quad \underline{t} = (t_i) \in R^N,$$

(by convention, $F_0 \in X$).

The class of hypercontractive random variables is closed under certain

algebraic operations and under weak limits of its distributions. The latter statement follows immediately from the definition of weak convergence of probability measures (see also proof of Proposition 2.2 in [6] for more details). The precise meaning of the first statement is contained in the following result (Krakowiak and Szulga [6]):

Theorem 2.1. Let $0 < q \leq p < \infty$, $\underline{\theta} = (\theta_i)$ be a sequence of independent p -integrable random variables, and F_0, \dots, F_k be finitely supported tetrahedral functions with values in X . If $\theta_j \in HC(p, q, X; s)$ for each $j=1, 2, \dots$ then

$$(2.1) \quad \left\| \sum_{k=1}^n \langle F_k; (\underline{s\theta})^k \rangle \right\|_p \leq \left\| \sum_{k=1}^n \langle F_k; (\underline{\theta})^k \rangle \right\|_q,$$

where $\underline{s\theta} = (s_i \theta_i)$.

In other words, the hypercontractivity is a hereditary feature under forming polynomials of several variables (not necessarily homogeneous). In particular, using linear forms, we obtain immediately the following statements:

COROLLARY 2.2 If (θ_i) is a sequence of i.i.d. random variables and each $\theta_i \in HC(p, q, X; s)$ then the space of all linear combinations of θ_i 's with X -valued coefficients consists of hypercontractive random variables, i.e.

$$\left\{ \sum_i x_i \theta_i : (x_i) \in X^N \right\} \in HC(p, q, X; s).$$

COROLLARY 2.3 If $\theta \in HC(p, q, X; s)$ and θ belongs to the domain of normal attraction of a random variable ξ then $\xi \in HC(p, q, X; s)$.

Therefore, by the central limit theorem argument, we obtain the following corollary

COROLLARY 2.4 (observed by Dr. Rama-Murthy). If $\theta \in HC(p, q, X; s)$ for some $s \neq 0$ and $E\theta^2 < \infty$ then $q > 1$.

Notice that if θ is a hypercontractive random variable then necessarily $|s| < 1$ and $E|x + y\theta|^q \geq 1$.

By homogeneity of the norms appearing in the definition of hypercontractivity it suffices to consider only vectors x with $\|x\| = 1$. Except a limited set of random variables, we may also assume that $\|y\| \leq 1$; the precise meaning of this remark is explained below.

LEMMA 2.5 Suppose that

$$\inf\{E\|x + y\theta\|^q; \|x\| = 1, \|y\| \geq 1\} > 1$$

and (1.2) is satisfied for all y with $\|y\| \leq 1$. Then $\theta \in HC(p, q, X; s)$ for some $s \neq 0$.

PROOF. Assume that $\|x\| = 1$. Let $t = \|y\| \geq 1$ and $q > 0$. Since

$$\lim_{t \rightarrow \infty} (\|x + y\theta\|_q^q - 1)/t = \|\theta\|_q^q$$

there is a positive constant c such that

$$\|1 + t\theta\|_q \geq 1 + ct \|\theta\|_q.$$

Let $r = \min(1, p)$. We solve the following inequality with respect to s , uniformly for $t \geq 1$,

$$(2.2) \quad \|x + sy\theta\|_p^q / \|x + y\theta\|_p^q \leq (1 + s^r t^r \|\theta\|_p^r)^{1/r} / (1 + ct \|\theta\|_q) \leq 1.$$

Define s_1 by

$$\begin{aligned} s_1 &= \inf_{t \geq 1} ((1 + ct \|\theta\|_q^r)^r - 1)^{1/r} / (t \|\theta\|_p) \\ &= ((1 + c \|\theta\|_q^r)^r - 1)^{1/r} / \|\theta\|_p > 0 \end{aligned}$$

and check that all s , $|s| \leq s_1$, satisfy (2.2), which completes the proof. \square

LEMMA 2.6. For any normed space X , any $q < 1$ and a symmetric random variable $\theta \in L^q$ we have

$$(2.3) \quad \delta_q(t) \stackrel{\text{df}}{=} \inf\{E\|x + y\theta\|^q; \|x\| = 1, \|y\| = t\} = E|1 + t\theta|^q.$$

PROOF. The inequality " \leq " is obvious.

Let $x, y \in X$ with $\|x\| = \|y\| = 1$ and x^* and y^* be functionals on X with norm 1 such that $\langle x^*, x \rangle = \langle y^*, y \rangle = 1$. Put $a = \langle x^*, y \rangle$ and $b = \langle y^*, x \rangle$. Then

$$\delta_q(t) \geq \inf\{E \max(|1 + a\theta|^q, |b + \theta|^q); 0 \leq a \leq 1, |b| \leq 1\}$$

Since, for a Bernoulli random variable σ independent of θ ,

$$E||x + y\theta||^q = E_\theta E_\sigma ||x + \sigma\theta y||^q,$$

it is enough to prove that for every $t \geq 0$

$$\inf \{E \max(|1 + a\sigma t|^q, |b + \sigma t|^q) : 0 \leq a \leq 1, |b| \leq 1\} \geq E|1 + t\sigma|^q.$$

Observe that, for $t \leq 1$ and $0 \leq a \leq 1$,

$$E|1 + a\sigma|^q \geq E|1 + t\sigma|^q \geq \inf_{|b| \leq 1} E|b + t\sigma|^q.$$

Therefore, for $t \leq 1$,

$$\begin{aligned} & \inf_a \inf_b E \max(|1 + a\sigma|^q, |b + t\sigma|^q) \\ & \geq \inf_a \inf_b \max(E|1 + a\sigma|^q, E|b + t\sigma|^q) \\ & \geq \max \left(\inf_a E|1 + a\sigma|^q, \inf_b E|b + t\sigma|^q \right) \\ & \geq \max \left(E|1 + t\sigma|^q, \inf_b E|b + t\sigma|^q \right) \\ & = E|1 + t\sigma|^q. \end{aligned}$$

Finally, for $t > 1$, we have

$$\inf_a E|1 + a\sigma|^q = 2^{q-1}.$$

Also, for a fixed $t > 1$, the function $b \rightarrow E|t + \sigma b|^q$ is decreasing for $b \in [0, 1]$, hence

$$\inf_b E|b + t\sigma|^q \geq E|1 + t\sigma|^q \geq 2^{q-1}. \quad \square$$

REMARK. In the case $q \geq 1$ the quantity $\delta_q(t)$, defined in (2.3), depends essentially on the geometry of the normed space X (cf. e.g. [14]).

If a random variable θ is integrable and $q \geq 1$ then the necessary and sufficient condition for hypercontractivity of θ is the finiteness of the function

$$(2.4) \quad V(s) = \sup_{t>0} f_p(st) / \{(1 + f_q(t) + \mu(t))^{p/q} - 1\},$$

where

$$\begin{aligned} f_p(t) &= E|t\theta|^p I\{|t\theta| > 2\}, \\ \mu(t) &= \inf \{E||x + t\theta y||^2 - 1\} I\{|t\theta| \leq 2\} : ||x|| = ||y|| = 1 \end{aligned}$$

(Krakowiak and Szulga [6]). Note that $\mu(t)$ may be replaced by any function $\mu'(t)$ with the same asymptotic behavior at 0. For example, if X is a Hilbert space then one may choose

$$\mu'(t) = E|t\theta|^2 I\{|t\theta| \leq 2\}$$

This criterion enables us to give an example of a non-hypercontractive random variable for certain parameters p and q , $1 < q < p < 2$.

EXAMPLE. Let θ be a symmetric random variable such that

$$N(x) = P(|\theta| > x) = x^{-\alpha} (\log x)^c, \quad x \geq e,$$

where $\alpha > 1$ and $c > 1$. Writing $f(t) \approx g(t)$ whenever $\lim_{t \rightarrow 0} f(t)/g(t) = \alpha_c$, $0 < |c| < \infty$, we derive the following asymptotic behavior of f_p and μ by a routine computation

$$f_p(t) \approx \begin{cases} N(1/t) & \text{if } p < \alpha, \\ N(1/t) \log(1/t) & \text{if } p = \alpha, \end{cases}$$

$$p(t) \approx \begin{cases} n(1/t) & \text{if } \alpha < 2, \\ t^2 & \text{if } \alpha \geq 2. \end{cases}$$

Therefore $V(s) = \infty$ for any $s \neq 0$ and for all p, q such that $1 < q < p = \alpha < 2$. In other words, for these parameters,

$$\theta \notin HC(p, q, R; s).$$

Unfortunately, the use of the criterion (2.4) is restricted to integrable random variables and to moments $p, q > 1$.

In the following sections a straightforward computation proves the hypercontractivity of all α -stable symmetric random variables, $0 < \alpha < 2$, in an arbitrary normed space, and, *a fortiori*, of all random variables whose distributions are weak limits of random α -stable multilinear forms (cf. discussion at the beginning of this section).

3. Hypercontractivity on the real line

Let ξ be a symmetric α -stable random variable. The aim of this section is to evaluate the absolute p^{th} moment of the transformed α -stable random variable $x + y\xi$. The exact integral formula

$$E|x + y\xi|^p = 2\Gamma(1+p)\sin(\pi p/2) \int_0^\infty (1 - \cos v \exp(-|vy|^\alpha)) v^{-p-1} dv$$

(cf. e.g. Zolotarev [15], p. 63) is too complex for our purposes, so we prefer to switch from ξ to an appropriate random variable from its domain of normal attraction.

THEOREM 3.1 *Let $0 < q < p < \alpha$. Then every α -stable symmetric random variable $\xi \in \text{HC}(p, q, \mathbb{R}; s)$ for some $s = s(p, q, \alpha) > 0$.*

PROOF. By virtue of Corollary 2.3 it is enough to prove that $\theta \in \text{HC}(p, q, \mathbb{R}; s)$ for some $s > 0$, where θ is the random variable belonging to the domain of normal attraction of ξ with the density

$$(3.1) \quad f(x) = \begin{cases} 0 & \text{if } |x| < 1, \\ \alpha/2 |x|^{-1-\alpha} & \text{if } |x| \geq 1. \end{cases}$$

Since L_p -norms are homogeneous and θ is symmetric, it suffices to evaluate the function $t \rightarrow E|1 + t\theta|^p$, $t \geq 0$. Define

$$\phi_p(u) = (|1 + u|^p + |1 - u|^p)/2.$$

We check that

$$(3.2) \quad \begin{aligned} E|1 + t\theta|^p &= \alpha t^\alpha \int_t^\infty \phi_p(u) u^{-1-\alpha} du \\ &= 1 + \alpha t^\alpha \int_t^\infty (\phi_p(u) - 1) u^{-1-\alpha} du \\ &= 1 + a_{p,\alpha} t^\alpha + g_{p,\alpha}(t), \end{aligned}$$

where

$$\begin{aligned} g_{p,\alpha}(t) &= \alpha t^\alpha \int_0^t (1 - \phi_p(u)) u^{-1-\alpha} du, \\ a_{p,\alpha} &= \alpha \int_0^\infty (\phi_p(u) - 1) u^{-1-\alpha} du \\ &= \alpha \int_0^1 \{(\phi_p(u) - 1) u^{-1-\alpha} + (\phi_p(u) u^{-p} - 1) u^{-1+\alpha}\} du \\ &= p/(\alpha - p)(1 - h(p, \alpha)). \end{aligned}$$

The two-parameter function h is given by the formula

$$h(p, \alpha) = \alpha(\alpha - p)/p \int_0^1 (1 - \phi_p(u))(u^{-1-\alpha} + u^{-1-p+\alpha}) du.$$

We claim that $h(p, \alpha) < 1$ for $0 < p < \alpha$. If $p > 1$ then $h(p, \alpha) < 0$ because $\phi_p(u) > 1$ for $p > 1$ and $u < 1$; if $p=1$ then $h(p, \alpha) = 0$ because $\phi_1(u) = 0$ for $u \leq 1$.

For the remaining p 's, observe that the function $p \rightarrow h(p, \alpha)$ is decreasing on the interval $(0, \min(1, \alpha))$ and

$$\lim_{p \rightarrow 0} h(p, \alpha) = 1 - (\alpha\pi/2) \cotan(\alpha\pi/2).$$

Indeed, integrating the series

$$1 - \phi_p(u) = - \sum_{k=1}^{\infty} \binom{p}{2k} u^{2k}, \quad 0 \leq u < 1,$$

term by term, we obtain the following representation of the function h

$$h(p, \alpha) = \alpha(\alpha - p) \sum_{k=1}^{\infty} (1-p) \dots (2k-1-p) ((2k-\alpha)^{-1} + (2k+\alpha-p)^{-1} / (2k)!.$$

A routine calculation shows that $p \rightarrow h(p, \alpha)$ is a decreasing function if $0 < p < \min(1, \alpha)$ and that

$$\lim_{p \rightarrow 0} h(p, \alpha) = \sum_{k=1}^{\infty} 2\alpha^2 / (4k^2 - \alpha^2) = 1 - (\alpha\pi/2) \cotan(\alpha\pi/2) < 1$$

(cf. e.g. Gradshteyn and Ryzhik [3]).

From the latter statement and from (3.2) we infer immediately that

$$(3.3) \quad E|1 + t\theta|^p > 1 \quad \text{for all } t > 0.$$

Moreover,

$$(3.4) \quad \lim_{t \rightarrow 0} g_{p, \alpha}(t)/t^2 = p(p-2)\alpha/(4-2\alpha).$$

Therefore there are positive constants $a = a_{p, \alpha}$ and $b = b_{p, \alpha}$ such that for all $t, 0 \leq t \leq 1$, we have

$$(3.5) \quad 1 + at^\alpha \leq E|1 + t\theta|^p \leq 1 + bt^\alpha.$$

The latter estimate yields the inequality

$$||1 + st\theta||_p \leq ||1 + t\theta||_q$$

valid for all $t, 0 \leq t \leq 1$, and all $s, |s| \leq s_0 \stackrel{\text{df}}{=} (b/a)^{1/\alpha}$.

We complete the proof by choosing $s = \min(s_0, s_1)$, where s_1 appears in Lemma (2.5).

In the following section we extend the above result to an arbitrary normed space.

4. Hypercontractivity in normed spaces.

The proof of the main result is based on some elementary inequalities.

LEMMA 4.1. Let $c > 0$ and $p > 0$. Define

$$f_p(v) = |(1+v)^p - pv|, \quad -1 \leq v \leq c.$$

Then

$$f_p(t) = \begin{cases} (1+v)^p - 1 - pv & \text{if } p \geq 1, \\ pv - (1+v)^p + 1 & \text{if } p \leq 1. \end{cases}$$

and there are constants $d_p, D_p > 0$ such that for all v , $-1 \leq v \leq c$,

$$d \leq f_p(v)/v^2 \leq D_p.$$

We omit the routine proof.

For fixed $x, y \in X$, $\|x\| = 1$, $\|y\| \leq c$ and a symmetric random variable η we define a stochastic process $V(s)$, $s \in [0, 1]$,

$$V(s) = \|x + s\eta y\| - 1$$

with values in $[-1, c]$.

LEMMA 4.2 For a symmetric integrable random variable η the following contractions hold true

$$(4.1) \quad V(s) \leq s V(1);$$

$$(4.2) \quad EV^2(s) \leq s^2 EV^2(1) + 4s EV(1).$$

PROOF. (4.1) follows from the convexity of the norm. Since

$$\begin{aligned} E\|x + s\eta y\|^2 &= E\|s(x + \eta y) + (1-s)x\|^2 \\ &\leq E(s\|x + \eta y\| + (1-s))^2 \\ &= E(sV(1) + 1)^2 \\ &= s^2 EV^2(1) + 2sEV(1) + 1, \end{aligned}$$

and

$$\begin{aligned}
E||x + s\eta y|| &= E||x - s\eta y|| \\
&= E||(1+s)x - s(x + \eta y)|| \\
&\geq (1+s) - sE||x + \eta y|| \\
&= 1 - sEV(1)
\end{aligned}$$

then

$$\begin{aligned}
EV^2(s) &= E(||x + s\eta y||^2 - 2||x + s\eta y|| + 1) \\
&\leq (s^2EV^2(1) + 2sEV(1) + 1) - (2 - 2sEV(1)) + 1 \\
&= s^2EV^2(1) + 4sEV(1). \quad \square
\end{aligned}$$

LEMMA 4.3 For every $a > c$,

$$(4.3) \quad (a - 1)^q \geq 1 + ((c - 1)^q - 1)/c^q a^q$$

if $c > 2$, and

$$(4.4) \quad (1 + a)^p \leq 1 + (1 + \frac{1}{c})^{(p-1)^+} a^p.$$

PROOF. Omitted. \square

LEMMA 4.4 There exists a constant $c > 2$ such that for every $x, y \in X$, with $||x|| = 1$ and $||y|| = t \leq 1$,

$$E(||x + \theta y||^q - 1)1\{|t\theta| \leq c\} > k_c t^\alpha.$$

PROOF. From Lemma 2.2 and Theorem 3.1 we infer that

$$\begin{aligned}
&E(||x + \theta y||^q - 1)1\{|t\theta| \leq c\} \\
&\geq E(|1 + \theta t|^q - 1)1\{|t\theta| \leq c\} \\
&= \alpha t^\alpha \int_t^c (E_\sigma |1 + \sigma u|^q - 1) u^{-\alpha-1} du \\
&> \alpha t^\alpha \int_0^c (E_\sigma |1 + \sigma u|^q - 1) u^{-\alpha-1} du \\
&= k_c t^\alpha. \quad \square
\end{aligned}$$

LEMMA 4.5. Let θ have the density (3.1). Then

$$\begin{aligned}
E|t\theta|^2 1\{|t\theta| \leq c\} &\leq \alpha c^{2-\alpha}/(2-\alpha)t^\alpha; \\
E|t\theta|^p 1\{|t\theta| > c\} &= \alpha c^{p-\alpha}/(\alpha-p)t^\alpha.
\end{aligned}$$

PROOF. Omitted. \square

Now we extend the main result of the previous section to an arbitrary normed space.

THEOREM 4.5. For every $p, q, 0 < q < p < \alpha < 2$, there is $s > 0$ such that every symmetric α -stable random variable $\xi \in HC(p, q, X; s)$, where X is an arbitrary normed space.

PROOF. It suffices to show that the random variable θ with density (3.1) (see the proof of Theorem 3.1) is hypercontractive. We may and do assume that $q < 1$ and $x, y \in X$ with $\|x\| = 1$, $\|y\| = t \leq 1$.

We set up directly a convenient decomposition

$$\begin{aligned} E\|x + s\theta y\|^p &= \\ 1 + E(\|x + s\theta y\|^p - 1)1\{|t\theta| \leq c\} + E(\|x + s\theta y\|^p - 1)1\{|t\theta| > c\} \\ &= 1 + A_p(t) + B_p(t), \end{aligned}$$

where c results from Lemma 4.4.

In order to find the needed upper estimate of $A_p(t)$ we apply Lemma 4.1, using notation and contractions from Lemma 4.2 with $\eta = \sigma\theta 1\{|t\theta| \leq c\}$, as follows

$$\begin{aligned} A_p(t) &= E(1 + V(s))^p - 1 \\ &= E(pV(s) + \operatorname{sgn}(p-1) f_p(V(s))) \\ &\leq ps EV(1) + (p-1)^+ D_p EV^2(s) \\ &\leq ps EV(1) + (p-1)^+ D_p (s^2 EV^2(1) + 4sEV(1)) \\ &= s(p + 4(p-1)^+)EV(1) + (p-1)^+ D_p s^2 EV^2(1) \\ &= s(p + 4(p-1)^+)/q (E(1 + V(1))^q - 1 + f_q(V(1)) + (p-1)^+ D_p s^2 EV^2(1)) \\ &\leq s(p + 4(p-1)^+)/q (E(1 + V(1))^q - 1) + s K_{p,q} EV^2(1) \\ &\leq s k_{p,q} E(\|x + \theta y\|^q - 1)1\{|t\theta| \leq c\} + s K_{p,q} E|t\theta|^2 1\{|t\theta| \leq c\}, \end{aligned}$$

where $k_{p,q}$ and $K_{p,q}$ are suitable positive constants.

By Lemma 4.3 and 4.5 there exists a constant $L_{p,q}$ such that

$$\begin{aligned} B_p(t) &\leq s^p L_{p,q} E(\|x + \theta y\|^q - 1)1\{|t\theta| > c\}, \\ K_{p,q} E|t\theta|^2 1\{|t\theta| \leq c\} &\leq L_{p,q} E(\|x + \theta y\|^q - 1)1\{|t\theta| > c\}. \end{aligned}$$

We complete the proof by selecting a number s such that

$$s^{\min(p,1)} L_{p,q} \leq p/(2q).$$

This yields the concluding inequality

$$\begin{aligned} (E||x + s\theta y||^p)^{1/p} &\leq (1 + p/q (E||x + \theta y||^q - 1))^{1/p} \\ &\leq E||x + \theta y||^q)^{1/q}. \end{aligned} \quad \square$$

COROLLARY 4.6. Let M be a symmetric α -stable measure, $0 < \alpha < 2$ independently scattered on the interval $[0, T]$, and let f_j , be function on $[0, T]^j$ with values in a Banach space X , integrable with respect to the product random measures M^j , $j=1, \dots, n$ (as defined in [7]). Put

$$(4.5) \quad \theta = \sum_{j=1}^n \int_{[0, T]^j} f_j dM^j.$$

Then θ is a hypercontractive random variable in X with parameters

p, q ($q < p < \alpha$) and s^n , where $s = s(p, q, \alpha)$ appears in Theorem 4.5. Note that an α -stable polynomial

$$\theta = \sum_{j=1}^n \sum_{k_1, \dots, k_j} a_j(k_1, \dots, k_j) \xi_{k_1} \dots \xi_{k_j}$$

is a special case of (4.5).

References

1. Bonami, A., Etude des coefficients de Fourier des fonctions de $L^p(G)$. *Annal. Inst. Fourier* 20, 335-402 (1970).
2. Borell, C., On polynomial chaos and integrability. *Probability Math. Statist.* 3, 191-203 (1984).
3. Gradshteyn, I.S. and Ryzhik, I.M., *Table of Integrals, Series and Products*. Academic Press 1980.
4. Gross, L., Logarithm Sobolev inequalities. *Amer. J. Math.* 97, 1061-1083 (1973).
5. Holley, R. and Stroock, D., Logarithm Sobolev inequalities and stochastic Ising model. *J. Statist. Physics.* 46, 5/6, 1159-1194 (1987).
6. Krakowiak, W. and Szulga, J., Hypercontraction principle and random multilinear forms. Wroclaw University Preprint #24 (1985) and CWRU Preprint series #85-34 (1985) (to appear in *Probability Theory and Related Fields*).
7. Krakowiak, W. and Szulga, J., A multiple stochastic integral with respect to a strictly p-stable random measure. CWRU Preprint Series #86-48 (to appear in *Ann. Probab.*)
8. Marcinkiewicz, J. and Zygmund, A., Sur les fonctions independantes. *Fundamenta Math.* 29, 60-90 (1937).
9. Nelson, E., A quartic interaction in two dimensions. In: *Mathematical theory of elementary particles*, 69-73, Cambridge, Mass., M.I.T. Press (1966).
10. Paley, R.E.A.C. and Zygmund, A., A note on analytic functions on the unit circle. *Proc. Cambridge. Phil. Soc.* 266-272 (1932).
11. Ritter, L.D., Convolution inequalities on finite cyclic groups and the pseudomeasure norm. *Proc. Amer. Math. Soc.* 91, 589-592 (1984).
12. Weissler, F.B., Logarithmic Sobolev inequalities and hypercontractive estimates on the circle. *J. Functional Anal.* 37, 218-234 (1980).
13. Wiener, N., The homogeneous chaos. *Amer. J. Math.* 60, 897-936 (1930).
14. Woyczynski, W.A., Geometry and martingales in Banach spaces. In *Lecture Notes in Math.* #472, pp. 229-276, Springer Verlag, Berlin-Heidelberg-New York 1975.
15. Zolotarev, V.M., *One-dimensional Stable Distributions*. AMS, Providence, Rhode Island, 1986 (translated from Russian).

END

DATE
FILMED

DEC.

1987